

Table 1.25  
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites<sup>a</sup> by Race/Ethnicity

## Females

	All Races		White		Black	
	Rate <sup>b</sup> 2013-2017	APC <sup>c</sup> 2008-2017	Rate <sup>b</sup> 2013-2017	APC <sup>c</sup> 2008-2017	Rate <sup>b</sup> 2013-2017	APC <sup>c</sup> 2008-2017
All Sites	418.3	-0.3*	432.0	-0.4*	390.7	-0.4*
Breast	128.5	0.2	131.3	0.0	124.8	0.3
Lung and Bronchus	48.6	-1.5*	51.5	-1.5*	43.8	-2.0*
Colon and Rectum	33.6	-2.2*	33.3	-2.0*	38.2	-3.0*
Corpus and Uterus, NOS	27.8	1.0*	28.3	0.7*	27.9	1.7*
Thyroid	23.1	0.8	24.3	0.6	14.3	1.5
Melanoma of the Skin	17.8	1.3*	21.7	1.4*	14.1	0.3
Non-Hodgkin Lymphoma	16.2	-0.8*	17.0	-0.9*	12.5	0.0
Pancreas	11.6	0.4*	11.7	-2.4*	12.2	-0.2
Ovary <sup>g</sup>	11.2	-2.1*	11.6	-0.4	12.0	0.3
Kidney and Renal Pelvis	11.1	0.3	11.6	0.4	9.1	-1.4*
Leukemia	11.0	-0.3	11.4	0.4	8.9	0.7
Urinary Bladder	8.6	-1.5*	9.4	-1.5*	8.7	-2.4*
Cervix Uteri	7.4	-0.7	7.2	-0.7	7.6	-2.1*
Oral Cavity and Pharynx	6.4	0.1	6.6	0.2	6.5	-0.4
Myeloma	5.7	0.6*	6.0	-0.6	5.4	0.8
	Asian/Pacific Islander		American Indian/Alaska Native <sup>d</sup>		Hispanic <sup>e</sup>	
	Rate <sup>b</sup> 2013-2017	APC <sup>c</sup> 2008-2017	Rate <sup>b</sup> 2013-2017	APC <sup>c</sup> 2008-2017	Rate <sup>b</sup> 2013-2017	APC <sup>c</sup> 2008-2017
All Sites	304.7	0.0	305.9	-0.4	339.5	0.2
Breast	102.9	1.2*	79.5	-0.5	99.1	0.6*
Lung and Bronchus	28.6	-0.2	37.9	0.0	29.2	-1.1*
Colon and Rectum	26.8	-3.0*	34.3	-1.0	24.8	-1.2*
Thyroid	22.7	0.8	19.9	-0.8	24.6	2.4*
Corpus and Uterus, NOS	21.7	1.9*	14.5	2.6*	22.3	2.3*
Non-Hodgkin Lymphoma	11.1	-0.6	13.7	-0.1	15.9	-0.2
Ovary <sup>g</sup>	9.5	-1.0	10.9	-1.2	11.7	1.1*
Pancreas	9.2	0.2	9.0	0.7	10.8	0.2
Stomach	8.0	-2.9*	8.4	-3.2	10.4	-0.9*
Liver & IBD <sup>f</sup>	7.2	-2.5*	7.9	0.4	9.2	-1.9*
Leukemia	6.5	-0.8	7.8	-0.6	9.1	-0.6
Cervix Uteri	6.5	-0.5	6.2	-1.1	8.6	-0.2
Kidney and Renal Pelvis	5.8	1.0	6.0	-3.4	8.1	1.6*
Oral Cavity and Pharynx	5.5	1.0	5.0	-0.2	5.7	0.3
Urinary Bladder	3.8	-1.6*	5.0	-	5.0	-1.9*

Source: SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

<sup>a</sup> Top 15 cancer sites selected based on 2013-2017 age-adjusted rates for the race/ethnic group.

<sup>b</sup> Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>c</sup> The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>d</sup> Rates for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

<sup>e</sup> Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

<sup>f</sup> IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

<sup>g</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

\* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.